



Belitung Nursing Journal

E-ISSN: 2477-4073 | P-ISSN: 2528-181X

Volume 8, Issue 3
May - June 2022

Edited by Assoc. Prof. Dr. Yupin Aungsuroch & Dr. Joko Gunawan

The Official Publication of
Belitung Raya Foundation
Department of Publication, Indonesia



Transitional care interventions to reduce emergency department visits in older adults: A systematic review

Latifah Jehloh^{ID}, Praneed Songwathana^{*ID}, and Wipa Sae-Sia^{ID}

Faculty of Nursing, Prince of Songkla University, Thailand



Abstract

Background: Preventable illnesses cause many emergency department visits in older adults, which can be minimized by implementing appropriate transitional care interventions. However, the most effective transitional care strategies for older adults are unknown.

Objective: To discover and consolidate transitional care interventions that can help older people avoid going to the emergency department.

Methods: From January 2011 to August 2021, PubMed, The Cochrane Library, CINAHL, Web of Science, ProQuest, and The JAMA Network were used to search. Two authors independently screened and selected papers, assessed the risk of bias, and extracted data into a standardized form in accordance with Cochrane guidelines. For the risk of bias in studies, the RevMan 5.4.1 program was utilized.

Results: Six randomized controlled trials, four non-randomized controlled trials, and three retrospective investigations were among the 13 studies examined. All studies evaluated emergency department visits but in different periods (ranging from 1-12 months after discharge) and with varying groups of baselines (pre-post intervention and between groups). The multi-component strategies, either pre or postdischarge phase using high-intensity care delivered within six months of discharge, were implemented in transitional care that had been shown to reduce emergency department visits in older adults.

Conclusion: To prevent emergency department visits by older patients, nurses should arrange for a high-intensity transitional care intervention that involves both pre-and postdischarge interventions. The effectiveness of the intervention in reducing emergency department visits in older adults is difficult to determine due to inter-study heterogeneity and poor methodological quality. There is a need for more evidence-based research with consistent and trustworthy effect assessments.

PROSPERO registration number: CRD42021261326

Keywords

emergency department visits; older adults; transitional care; interventions; systematic review; discharge planning; nursing

*Corresponding author:

Assoc. Prof. Dr. Praneed Songwathana
Faculty of Nursing, Prince of Songkla
University, Songkhla 90110, Thailand
Email: praneed.s@psu.ac.th

Article info:

Received: 22 March 2022

Revised: 24 April 2022

Accepted: 30 May 2022



This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License, which allows others to remix, tweak, and build upon the work non-commercially as long as the original work is properly cited. The new creations are not necessarily licensed under the identical terms.

E-ISSN: 2477-4073 | P-ISSN: 2528-181X

Background

The number of aging persons has been increasing dramatically worldwide over the past few decades (Feliciano et al., 2022). The aging population is linked to an increased burden of non-communicable disease disorders, all-function organ dysfunction (Amarya et al., 2018), poor health status (McPake & Mahal, 2017), and complex medical needs (Chandra et al., 2015), which necessitate long-term care (McPake & Mahal, 2017). Age, functioning, polypharmacy, pharmacotherapy complexity, inappropriate drugs used for older persons, and multimorbidity contribute to the high demand for care. Return visits to the Emergency Department (ED) following hospital discharge can be linked to such factors

(Deschodt et al., 2015; Naseer et al., 2020; Salvi et al., 2017; Schoonover et al., 2014).

ED visits are common among older adults (Pham et al., 2009). Previous research has shown that 10-26 percent of older persons attend the emergency room (de Gelder et al., 2018; Lowthian et al., 2016). Aging patients made more than five ED visits per year, accounting for 15% of all visits (Ukkonen et al., 2019). Furthermore, dyspnea from COPD was the most common chief complaint expressed by the elderly with disease-related visits to the ED (Paksopis et al., 2019; Song et al., 2016). A previous study reported that nearly 55.2 percent of all ED visits were older adults with COPD (Hasegawa et al., 2014). A high rate of visiting ED among persons with COPD occurred after hospital discharge, mostly due to the symptom exacerbation (Paksopis et al., 2019; Song

et al., 2016). Therefore, it is necessary to prevent acute exacerbation in COPD patients at home, which requires transitional care to treat and stabilize patients' symptoms to return to their normal lives outside the hospital. Optimal transitional care from hospital to home in aged care is quality in health care standards and indicators as recommended in the research field of transitional care (Allen et al., 2014). Additionally, the use of emergency services after being discharged from the hospital can be a predictor that the transitional care process is inadequate (Santos et al., 2019).

Transitional care is a broad term for medical approaches that facilitate the safe and timely transition of elderly patients across care levels and settings (Naylor & Keating, 2008) from pre-hospital discharge and prompt follow-up following hospital discharge (Naylor et al., 2011). Transitional care for elderly patients is essential to minimize medication errors, adverse drug events, lack of timely follow-up care management, and unnecessary ED visits (Kessler et al., 2013). In addition, transitional care is required to avoid discontinuity of treatment and adverse effects on older patients. On the other hand, the poor transitional care led to increased emergency treatment (Son & You, 2015). Therefore, Transitional Care Interventions (TCIs) in the older population are interesting to prevent adverse events during the transition and reduce ED visits.

Few systematic reviews of studies are available on what is the best transitional care intervention to reduce ED visits in older individuals. In 2020, similar research released a systematic review on the impact of transitional care interventions on hospital readmissions in elderly medical patients (Rasmussen et al., 2021). However, the study's primary outcome is hospital readmission, and the findings are equivocal, indicating that there is no evidence to advocate a specific intervention. As a result, more knowledge on interventions that minimize ED visits by older adults is needed to help health care workers, particularly nurses, identify and implement effective transitional care interventions to reduce the ED visit in these populations.

Methods

Design

A systematic review design was employed. Prior to data collection, this review was entered into the PROSPERO database (CRD42021261326). It follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (PRISMA).

Eligibility Criteria

Studies that focused on interventions towards older adult patients were considered. All studies, including older adults discharged from a general medical ward or emergency department using the PICO framework, were included (Table 1). In addition, interventions aimed at people in the transitional phase were studied, and the impact on ED visits was calculated. The interventions were required to involve either pre- or postdischarge components by either the provider or the patient management, with the number of ED visits being the primary measure. Randomized controlled trials (RCTs), quasi-experimental designs, retrospective studies, cohort studies, and case-controlled studies were required. Studies were

omitted if the target population was under 60 years old or if the result was not an ED visit.

Table 1 PICO format

PICO format	
Population	Older patients
Intervention	Transitional care interventions
Comparison	Usual care
Outcomes	ED visits

Data Sources

From January 2011 to August 2021, this study examined the following bibliographic databases to find relevant studies: PubMed, The Cochrane Library, Cumulative Index to Nursing and Allied Health Literature (CINAHL), Web of Science, ProQuest, and The JAMA Network.

Search Strategy

Only studies in English published during the year 2011-2021 were included in the searches. The bibliographic searches took place between 15 July and 6 August 2021. Searching terms applied to search articles were based on MESH terms "transitional care, discharged intervention, emergency room visit, older adults". Keyword and inclusion criteria used for searching in each database are shown in Table 2.

Study Selection

First, the first author assessed titles for their possible relevance to the population and outcome (LJ). Second, two authors independently reviewed titles and abstracts to see if they were intervention-worthy (LJ and PS). Third, two authors separately assessed the entire material. In the event of a disagreement, a third author (WS) was validated and discussed.

Quality Appraisal

The Joanna Briggs Institute (JBI)'s critical appraisal techniques were used to evaluate the trustworthiness, relevance, and findings of published publications (Joanna Briggs Institute, 2020).

Data Extraction

The Cochrane Data Extraction Form was altered to accommodate the current patient group and intervention type (Cochrane, 2022). Two researchers independently extracted data from the listed studies. Author, year of publication, nation, study design, setting, participants, study size, outcomes, follow-up period, and impact of intervention in numbers and/or percent were among the study features and results extracted. Only information about the 'ED visit' outcome was extracted and evaluated.

Assessment of Risk of Bias

Two reviewers (LJ and PS) separately rated the methodological quality. The papers were assessed using the risk of bias criteria according to Cochrane Effective Practice and Organisation of Care (EPoC) (2017). Allocation sequence generation, concealment of allocation, similar baseline outcome measurements, similar baseline characteristics, incomplete outcome data, participant blinding, outcome assessor blinding, contamination protection, selective

outcome reporting, and other potential sources of bias in studies were evaluated by two reviewers. Validation and determination of whether the requirements were completed were done through discussion or consultation with a third researcher (WS). According to Davey et al. (2013), reviewers classified each study as 'Low' if all criteria were rated as 'Low risk,' 'Moderate' if one or two criteria were rated as 'Unclear' or 'High risk,' and 'High' if more than two criteria were rated as 'Unclear' or 'High risk'.

Data Synthesis and Analysis

Methodological quality, intervention characteristics, outcomes, statistical significance, and the direction of observed effects were all used to characterize the research findings. One primary reviewer (LJ) reviewed each study and recorded all of the characteristics of the therapy investigated. Individual research was looked at further to check if they utilized the

same or different terminology to describe the same elements. A set of core elements was created as a result of this method. After then, each individual intervention was evaluated once more to see if it fit into one of the categories or not. The prior processes were double-checked by a second reviewer (PS), who was a student's supervisor. Summaries of intervention effects were computed for each trial, when possible, using risk ratios (RR) for binary outcomes and standardized mean differences (SMD) for continuous outcomes. This review attempts to do meta-analyses to determine the effectiveness of transitional care intervention by using RevMan V 5.4.1 software (<https://training.cochrane.org/online-learning/core-software-cochrane-reviews/revman/revman-5-download/download-and-installation>). The meta-analysis process could not be completed because the included studies had poor methodological studies, discrepancies in follow-up measurement, and multi-component characteristics of the intervention.

Table 2 Keywords and inclusion criteria used for searching in each database

Databases	Main search	Limit	Search found (N)	Meet criteria (N)	Reasons for exclusion
ProQuest	("transitional care" OR "discharge planning") AND ("intervention") AND ("emergency department visit" OR "emergency room visit") AND ("older adults" OR "elderly" OR "aged")	<ul style="list-style-type: none"> English Full-text Year 2011-2021 Scholarly journals 	869	22	Outcome and population are not matched; not intervention studies; not transitional care interventions; systematic review
CINAHL	("transitional care" OR "care transition" OR "discharge planning") AND ("intervention" OR "program") AND ("emergency department visit" OR "emergency room visit") AND ("older adults" OR "elderly" OR "geriatrics" OR "aging" OR "aged")	<ul style="list-style-type: none"> English 2011-2021 Full-text Academic journal Age: older adults 	135	16	The outcome is not matched; not intervention studies
Web of Science	("transitional care" OR "discharge planning") AND ("intervention" OR "program") AND ("emergency department visit" OR "emergency room visit") AND ("older adults" OR "elderly" OR "geriatrics" OR "aging" OR "aged")	<ul style="list-style-type: none"> English Full-text Year 2011-2021 	61	14	Not transitional care interventions; outcome is not matched; not intervention studies
PubMed	("transitional care") OR ("transition of care") OR ("discharge planning") AND ("intervention" OR "program") AND ("emergency department visit" OR "emergency room visit") OR ("emergency department attendance") OR ("emergency room attendance") AND ("older adults") OR ("elderly") OR ("geriatrics") OR ("aging") OR ("aged")	<ul style="list-style-type: none"> Year 2011-2021 Free full-text Human English Aged: 65+ years 	127	15	The outcome is not matched
The Cochrane Library	("transitional care" OR "discharge planning") AND ("intervention") AND ("emergency department visit" OR "emergency room visit") AND ("older adults" OR "elderly")	<ul style="list-style-type: none"> Trials Year 2011-2021 	34	3	The outcome is not matched; not transitional care interventions
The JAMA Network	("transitional care" OR "discharge planning") AND ("intervention" OR "program" OR "strategies") AND ("emergency department visit" OR "emergency room visit") AND ("older adults" OR "elderly")	<ul style="list-style-type: none"> Geriatrics Research Free Year 2011-2021 	60	4	The outcome is not matched; not transitional care interventions

Results

Study Selection

A total of 1152 records were found, and 1085 records were screened by title and abstract after duplicates were removed. A total of 59 records were examined for full-text review, with 13 meeting the eligibility requirements. Participants' age, population, research design, or ED visit were not reported as outcomes in the 46 remaining entries, as shown in [Figure 1](#).




Figure 1 PRISMA flow diagram

Study Characteristics

The 13 trials included in the study contained 13 different therapies and outcome evaluations that measured ED visits at various time intervals. [Table 3](#) lists the study's features. Six of the studies included were RCTs ([Enguidanos et al., 2012](#); [Legrain et al., 2011](#); [Liang et al., 2021](#); [Mi et al., 2018](#); [Occelli et al., 2016](#); [Schapira et al., 2021](#)), four were non-randomized controlled trials (NRCTs) ([Heeren et al., 2019](#); [Lee et al., 2018](#); [Low et al., 2015](#); [Robinson et al., 2021](#)) and three were retrospective studies ([Kim et al., 2021](#); [Sabir et al., 2019](#); [Wee et al., 2014](#)). Individual study sample sizes ranged from 199 to 4835 people. Within 1–12 months of discharge, outcome evaluations were undertaken. Aside from ED visits, the 13 studies looked at hospital readmission ([Heeren et al., 2019](#); [Kim et al., 2021](#); [Liang et al., 2021](#); [Low et al., 2015](#); [Sabir et al., 2019](#); [Schapira et al., 2021](#); [Wee et al., 2014](#)), the mortality rate ([Enguidanos et al., 2012](#); [Heeren et al., 2019](#); [Liang et al., 2021](#)), functional decline ([Heeren et al., 2019](#)), health outcome ([Lee et al., 2018](#)), the impact of polypharmacy ([Sabir et al., 2019](#)), health care cost ([Kim et al., 2021](#)), self-efficacy ([Enguidanos et al., 2012](#)), and patient's satisfaction

([Enguidanos et al., 2012](#)). Three studies were carried out in the United States ([Enguidanos et al., 2012](#); [Kim et al., 2021](#); [Mi et al., 2018](#)), two in France ([Legrain et al., 2011](#); [Occelli et al., 2016](#)), two in Singapore ([Low et al., 2015](#); [Wee et al., 2014](#)), one in Argentina ([Schapira et al., 2021](#)), one in the New Zealand ([Robinson et al., 2021](#)), one in Belgium ([Heeren et al., 2019](#)), one in the Hongkong ([Lee et al., 2018](#)), one in the UK ([Sabir et al., 2019](#)), and one in Taiwan ([Liang et al., 2021](#)). Between 2011 and 2021, the research was published.

Study Population

This review included roughly 16,105 patients, with 6750 in the intervention groups and 9,355 in the control groups. The participants in the intervention groups were on average 82 years old (range 77.7–85.9), while those in the control groups were on average 82 years old (range 77.9–86.4).

Transitional Care Interventions (TCIs)

The strength of each TCI is described in [Table 4](#). [Verhaegh et al. \(2014\)](#) came up with the idea of calculating intervention intensity. Only one study was classified as low intensity ([Sabir et al., 2019](#)). The majority of included studies have a high intensity ([Enguidanos et al., 2012](#); [Heeren et al., 2019](#); [Kim et al., 2021](#); [Lee et al., 2018](#); [Legrain et al., 2011](#); [Liang et al., 2021](#); [Low et al., 2015](#); [Mi et al., 2018](#); [Occelli et al., 2016](#); [Robinson et al., 2021](#); [Schapira et al., 2021](#); [Wee et al., 2014](#)). The vast majority of high-intensity interventions had a beneficial influence on ED visits, but one low-intensity intervention also had a good impact. The effects of high-intensity interventions were bigger and statistically significant.

The ED visits in these studies were monitored over a period ranging from one month to twelve months after hospital discharge. Eight studies looked at ED visits six months after discharge ([Enguidanos et al., 2012](#); [Kim et al., 2021](#); [Lee et al., 2018](#); [Legrain et al., 2011](#); [Liang et al., 2021](#); [Low et al., 2015](#); [Sabir et al., 2019](#); [Schapira et al., 2021](#)), and seven studies looked at three months after discharge ([Heeren et al., 2019](#); [Kim et al., 2021](#); [Lee et al., 2018](#); [Legrain et al., 2011](#); [Liang et al., 2021](#); [Low et al., 2015](#); [Wee et al., 2014](#)), five studies observed at one month after discharge ([Heeren et al., 2019](#); [Kim et al., 2021](#); [Mi et al., 2018](#); [Occelli et al., 2016](#); [Wee et al., 2014](#)). The rest (just one trial) was assessed either two months after discharge ([Kim et al., 2021](#)) or one year after discharge ([Robinson et al., 2021](#)). At one month, two months, and three months after discharge, a high-intensity TCI intervention would considerably reduce ED visits for older adults. However, the results on the intensity of the intervention are ambiguous six months after discharge because two studies with high-intensity treatments found no significant differences between the two groups in any of the ED visits ([Lee et al., 2018](#); [Legrain et al., 2011](#)), while one study with a low-intensity intervention found a beneficial effect on ED visits ([Sabir et al., 2019](#)). Furthermore, 12-month high-intensity intervention has a beneficial influence on ED visits, according to one study ([Robinson et al., 2021](#)).

Table 3 Study characteristics of pre-discharge – post-discharge interventions

Author (year)	Country setting	Design	Intervention	Study population	Outcomes	ED visit measured time	ED visit rate (IG)	ED visit rate (CG)	RR/OR (95%CI)
Schapira et al. (2021)	A tertiary care university hospital, Buenos Aires, Argentina	Single-blinded RCT	Predischarge: 1. A comprehensive geriatric assessment 2. Tailored recommendation 3. Transition of care Postdischarge: 1. A health and social care counselor	Adults > 75 Total 240 (IG=120, CG=120)	Hospital readmission; ED visit; Mortality rate	6 months after D/C	N = 120 n = 52 43.3%	N = 120 n = 72 60%	RR = 0.722 (0.562- 0.929)
Robinson et al. (2021)	Auckland, New Zealand	A non- RCT	Predischarge: 1. A comprehensive geriatric assessment 2. Transition of care 3. Care planning 4. Self-management education Postdischarge: 1. Follow-ups	People aged >75 IG =1085 CG = 3750	ED visit; Mortality rate	12 months after D/C	N = 1085 n = 56 0.05%	N = 3750 n = 420 0.11%	OR = -0.07 (-0.10-0.04) p <0.001
Low et al. (2015)	Largest tertiary hospital, Singapore	A quasi-experimental study using a pre-post design	Predischarge: 1. Care plan Postdischarge: 1. Care plan 2. A comprehensive need assessment	259 elderly aged> 60 years	Hospital readmission; ED visit	3 months after D/C 6 months after D/C	N = 259 Pre 410 Post 217 47.1% reduction N = 259 Pre 625 Post 324 48.2% reduction	-	NR
Heeren et al. (2019)	University Hospitals, Leuven, Belgium	A quasi-experimental study (sequential design with two cohorts)	Predischarge: 1. Care plan Postdischarge: 2. Follow-ups	Patients aged >70 years CG=768 IG= 857	ED visits; hospitalization rate; functional decline; and mortality.	1 month after D/C 3 months after D/C	N = 857 n = 112 13.1% N = 857 n = 205 23.9%	N = 768 n = 93 12.1% N = 768 n = 170 22.1%	NR
Lee et al. (2018)	A community in Hong Kong	A matched-control quasi-experimental design	Postdischarge: 1. Rehabilitation 2. Cognitive and memory training	122 pairs of older adults and their caregivers	Health outcome; ED visits	3 months after D/C 6 months after D/C	N = 61 -0.66 ±1.03 N = 61 -0.49 ±0.96	N = 61 0.05 ±0.76 N = 61 0.07 ±0.68	NR
Sabir et al. (2019)	Leeds Teaching Hospitals NHS Trust (LTHT), UK	Retrospectively collected	Predischarge: 1. A patient information leaflet and a web-page Postdischarge: 1. Refer information/ an outcome to a community pharmacist	621 patients aged 65 and over	Hospital readmission; ED visit; Impact of polypharmacy	6 months prior to D/C 6 months after D/C	N = 621 Pre 214 34.13% Post 171 27.27%	NC	NR
Liang et al. (2021)	National Yang-Ming University Hospital, Taiwan	RCT	Predischarge: 1. Provided wireless transmission devices Postdischarge: 1. Telemonitoring through wireless transmission devices 2. Home visit	IG = 100 CG= 100	Hospital readmission; ED visit; Mortality rate	6 months after D/C	N = 100 n = 12 12%	N = 100 n = 26 26%	OR =0.388 (0.183-0.822) P = 0.013

Table 3 (Cont.)

Kim et al. (2021)	A primary care internal medicine residency clinic, USA	Retrospective study	Predischarge: 1. Medication counseling Postdischarge: 1. Phone visit 2. Face to face teach back 3. Health education	65 older patients	Hospitalization n; ED visits; Health care cost	1 month after D/C 2 months after D/C 3 months after D/C 6 months after D/C	<i>N</i> = 49 <i>n</i> = 2 <i>n</i> = 2 <i>n</i> = 6 <i>n</i> = 25	<i>N</i> = 52 <i>n</i> = 12 <i>n</i> = 10 <i>n</i> = 17 <i>n</i> = 40	NR
Enguidanos et al. (2012)	USA	RCT	Postdischarge: 1. Home visit 2. Phone call visit	Older adults (<i>N</i> = 199)	Self-efficacy; Patient's satisfaction; ED visit	6 months after D/C	<i>N</i> = 100 <i>n</i> = 0.5 1.2%	<i>N</i> = 99 <i>n</i> = 0.99 2.5%	NR
Legrain et al. (2011)	Six acute geriatric units (AGUs) in Paris and its surroundings.	Open-label, six-month prospective, randomized (Zelen design), parallel-group trial	Predischarge: 1. Comprehensive chronic medication review 2. Self-management education 3. Transition of care communication with outpatient health professionals	All 665 Aged > 70 IG = 317 CG = 348	ED visit	3 months after D/C 6 months after D/C	<i>N</i> = 317 <i>n</i> = 11 3.8% <i>n</i> = 19 6.0%	<i>N</i> = 348 <i>n</i> = 12 3.4% <i>n</i> = 22 6.3%	NR
Mi et al. (2018)	Rochester, NY and Madison, USA	Single-blinded RCT	Postdischarge: 1. Home visit 2. Phone call visit	2400 patients over the age of 60 were enrolled.	ED visit	1 month after D/C	NR	NR	NR
Occelli et al. (2016)	The Rhône-Alpes region of France	A stepped-wedge cluster randomized experiment using a multicentre-design	Predischarge: 1. Discharge care plan Postdischarge: 1. Follow-ups	630 patients aged > 75	ED visit	30 days after D/C	NR	NR	NR
Wee et al. (2014)	Five hospitals in Singapore	A retrospective cohort study	Predischarge: 1. Coaching 2. Care plan Postdischarge: 1. Follow-ups	IG = 4132 CG = 4132 aged > 65	Hospitalization n; ED visits	1 month after D/C 6 months after D/C	<i>N</i> = 4132 <i>n</i> = 992 19.3% <i>n</i> = 3801 46.3%	<i>N</i> = 4132 <i>n</i> = 1240 32% <i>n</i> = 4545 57.9%	<i>OR</i> = 0.81(0.72-0.90) <i>p</i> < .001 0.90 (0.82-0.99) <i>p</i> = .03

IG= intervention group, CG= control group

Predischarge: Transitional care interventions that are implemented before discharge

Postdischarge: Transitional care interventions are those that are undertaken after a patient is discharged

Transitional care intervention: Interventions assist elderly persons in regaining their independence following a hospital stay. It is provided either before or after the discharge of older patients

ED visit: The ED visits after discharge from hospital/ED in measured time (1,2,3,6 and 12 months after discharge)

The effect of the intervention on the outcome: reduced ED visit

NR = not report

Predischarge Components (TCIs before discharge from hospital)

A comprehensive geriatric assessment, tailored recommendations to minimize geriatric syndromes, planned transition of care, care plan, self-management education, a patient information leaflet, and a web page, wireless transmission devices, medication counseling, comprehensive chronic medication review, transition-of-care communication with outpatient health professionals, discharge care planning, and coaching were all included in predischarge components across studies. The most widely used TCIs of predischarge intervention in these articles that were helpful in reducing ED visits among older persons were care planning, the transition of care, and self-management education.

Post Discharge Components (TCIs after discharge from hospital)

A health and social care counselor, regular follow-up, a comprehensive needs assessment, an individualized care plan, rehabilitation, cognitive and memory training, information/outcome referral to community pharmacist, telemonitoring through wireless transmission devices, home visit, phone visit, face-to-face demonstration, and health education were all included in the postdischarge components. Follow-ups, home visits, and phone calls were the most commonly employed TCIs of postdischarge intervention in these papers that were helpful in reducing ED visits among older persons.

Table 4 The intensity of the intervention

Author	Assessment at admission	Self-management education	Caregiver involvement	Discharge planning	Care coordination by the nurse	Primary care provider communication	Home visit within three days	Number of the home visit	Telephone follow-up	Number of telephone follow-up	The total duration of intervention	The total score of intervention intensity
Schapira et al. (2021)	/	/	/	/	/	/	/				3	10
Robinson et al. (2021)	/	/	/	/	/	/			1	1	3	11
Low et al. (2015)	/	/	/	/	/	/	/		1		3	11
Heeren et al. (2019)	/	/	/	/	/	/			1	2	3	11
Lee et al. (2018)	/	/	/	/	/	/	/	1			3	11
Sabir et al. (2019)	/			/							3	5
Liang et al. (2021)	/		/		/	/	/	1			3	9
Kim et al. (2021)		/			/	/	/		1	2	3	10
Enguidanos et al. (2012)	/	/	/	/	/	/	/	3	1	2	3	16
Legrain et al. (2011)	/	/	/	/	/	/					3	9
Mi et al. (2018)	/			/		/	/	1	1	2	1	9
Occelli et al. (2016)	/			/	/	/	/	2	1	2	1	11
Wee et al. (2014)	/	/		/		/		2	1	2	3	12

Note: / symbol for yes

Risk of Bias in Included Studies

Figure 2 and **Figure 3** show the overall judgment scores for each risk of bias item. Ten of the thirteen trials (76.9%) showed a high risk of bias owing to insufficient randomization, allocation concealment, and blinding (Enguidanos et al., 2012; Heeren et al., 2019; Kim et al., 2021; Lee et al., 2018; Liang et al., 2021; Low et al., 2015; Robinson et al., 2021; Sabir et al., 2019; Schapira et al., 2021; Wee et al., 2014). There was a moderate probability of bias in three studies (23.1%) (Legrain et al., 2011; Mi et al., 2018; Occelli et al., 2016). The overall risk of bias was considerable in the majority of the included trials due to different shortcomings in the randomization procedure in older persons. Older persons are commonly underrepresented in RCTs (Broekhuizen et al., 2015). As a result, clinical trial participants rarely represent older patients in general medical practice, putting RCTs' external validity in the older patient population in jeopardy (Kennedy-Martin et al., 2015). In addition, evidence-based treatment has limited application in this population because older people are usually excluded from clinical trials (Fougère et al., 2016). As a result, the scope of the RCT investigation among these people is limited. Therefore, investigations with a high chance of bias are still required.




Figure 2 Risk of bias graph: Review authors' decisions about each risk of bias item using RevMan 5.4.1

Synthesis of Results

When it came to the impact on the outcome, all studies (100%) showed a positive impact on ED visits 1, 2, and 3 months following hospital discharge, but not on TCIs conducted after six months (Lee et al., 2018; Legrain et al., 2011). Only three of the thirteen research studies (Liang et al., 2021; Robinson et al., 2021; Wee et al., 2014) provided an impact size estimate. Odds ratios (ORs) ranged from -0.07 to 0.90. One study provides an impact size by using a Risk Ratio (RR) (Schapira et al., 2021) ranging from 0.562 to 0.929. Because only four of the 13 studies provided the risk ratio and odds ratios, it was unable to calculate RRs and ORs and their 95 percent confidence intervals for those studies.

Meta-Analysis

Because most of the included studies had a significant risk of bias and differences in follow-up measurement intervals and multi-component characteristics, only three studies mentioned odd ratios, and meta-analytic pooling for impact estimations of effect size was problematic.




Figure 3 Risk of bias summary using RevMan 5.4.1

Discussion

According to our data, the majority of interventions in the transitional period between hospital and home appeared to reduce the rate of ED visits among older patients discharged from the ED and a medical ward. TCIs undertaken more than six months after discharge from the hospital, on the other hand, gave inconclusive results. There are several possible causes for these inconsistencies, as discussed below.

Our findings indicate that the intensity of interventions has an impact on the rate of ED visits. High-intensity interventions had a greater impact than low-intensity interventions in general. This is consistent with prior findings (Verhaegh et al.,

2014), which show that intervention components that appear to be the same across trials might have different characteristics. The varied influence on ED visits could be due to differences in intervention components between trials. A cross-study intervention component analysis to establish which components have a positive influence on ED visits is impossible. A full geriatric assessment, need assessment, discharge planning, caregiver involvement, self-management education, coaching, home visits, and telephone call visits were all components of interventions that had a positive influence on ED visits. Furthermore, the timing of the outcome evaluation influences the chance of a preventive ED visit. The findings of this study imply that interventions with high intensity and duration at 1, 2, and 3 months after discharge from the hospital are beneficial.

Several of the studies included did not account for potential confounders such as growing age, gender, psychological problems, poor general health (Sheikh, 2019), previous experience history, living in a rural location, low income, polypharmacy, comorbidity, and dementia (Dufour et al., 2019). Internal validity and the influence on ED visit rates may have been harmed as a result of these factors. Furthermore, randomized controlled trials (RCTs) made up less than half of the research considered (Enguidanos et al., 2012; Legrain et al., 2011; Liang et al., 2021; Mi et al., 2018; Occelli et al., 2016; Schapira et al., 2021). As a result, ED visit rates in these studies may be skewed.

The bulk of the included studies had poor methodological quality, indicating a substantial risk of bias. A low measure grade can show poor methodological quality and inadequate reporting techniques. The latter makes determining the genuine quality and the possibility of bias challenging. If the low ranking is due to insufficient methodological reporting, the effects may be unaffected. If the poor rating is due to methodological issues, however, it is likely to underestimate the true impact of the interventions. The research design and confounders component receive the lowest score. Because older adults are frequently omitted from clinical trials, RCTs and consideration of confounder variables are not always possible in older adult population research (Fougère et al., 2016). Furthermore, no matter how many variables the researcher controls for, residual confounding will exist, maybe due to unknown and unmeasured causes (Skelly et al., 2012). As a result, the low rating of methodological quality in study design and managing confounder variables is thought to be linked to the numerous constraints of doing research with the elderly.

Study Limitation

This review tries to synthesize the existing transitional care program between hospital and home, with the goal of reducing postdischarge ED visits for older patients. Although the analysis highlights the benefits of transitional care interventions in improving ED visits in these populations, it also identifies some potential limits. Small studies were considered; searches were limited to English-language literature, and studies in this population were few. We also didn't have enough research to do a meta-analysis because only four studies examined the transitional care intervention's effect size. As a result, the most effective transitional care treatments for reducing ED in this population are unclear.

Implications of this Study for Nursing Practice

In terms of clinical implications, nurses should plan for a high-intensity transitional care intervention that includes both pre- and postdischarge interventions in order to reduce ED visits by older patients. Future meta-analyses will be required to determine the efficacy of transitional care interventions.

Conclusion

The majority of TCIs have a beneficial influence on ED visit rates among older patients, with the greatest benefit occurring less than six months following discharge. As a result, we feel that the present evidence supports the recommendation of pre-discharge and postdischarge transitional care. However, there is insufficient data to propose a specific intervention. The major finding is that high-intensity therapies had a pronounced favorable impact within less than six months of discharge. However, this should be seen in the context of the fact that only 13 research matched the inclusion criteria, and the methodological quality approach assigned a low level of certainty to the evidence.

Declaration of Conflicting Interest

The researchers declare that they have no competing interests.

Funding

The Ministry of Higher Education, Science, Research and Innovation (MHESRI), Thailand, and the Research Center for Prevention and Care for People in Emergency and Trauma, Prince of Songkla University, Thailand.

Acknowledgment

The researchers would like to express their sincere thanks to the Ministry of Higher Education, Science, Research and Innovation (MHESRI), Thailand, and the Research Center for Prevention and Care for People in Emergency and Trauma, Prince of Songkla University, Thailand, for their financial support. In addition, the researchers are grateful to Mr. Alan Frederick Geater from the Faculty of Medicine, PSU, for his editorial assistance.

Authors' Contributions

LJ was in charge of the study's design, as well as writing and revising the manuscript. LJ and PS examined the data, authored the manuscript, and revised it. The study was developed by WS, who also wrote and revised the report. The final version of the paper to be published was approved by all authors.

Authors' Biographies

Latifah Jehloh is a PhD Candidate at the Faculty of Nursing, Prince of Songkla University, Thailand.

Praneed Songwathana, PhD, RN is an Associate Professor, Faculty of Nursing, Prince of Songkla University, Thailand.

Wipa Sae-Sia, PhD, RN is an Associate Professor, Faculty of Nursing, Prince of Songkla University, Thailand.

Data Availability

The datasets generated and analyzed during the current study are available from the corresponding author on reasonable request.

References

Allen, J., Hutchinson, A. M., Brown, R., & Livingston, P. M. (2014). Quality care outcomes following transitional care interventions for older people from hospital to home: A systematic review. *BMC Health Services Research*, 14(1), 1-18. <https://doi.org/10.1186/1472-6963-14-346>

- Amarya, S., Singh, K., & Sabharwal, M. (2018). Ageing process and physiological changes. In G. D'Onofrio, A. Greco, & D. Sancarolo (Eds.), *Gerontology*. IntechOpen. <https://doi.org/10.5772/intechopen.76249>
- Broekhuizen, K., Pothof, A., de Craen, A. J. M., & Mooijaart, S. P. (2015). Characteristics of randomized controlled trials designed for elderly: A systematic review. *PloS One*, 10(5), e0126709. <https://doi.org/10.1371/journal.pone.0126709>
- Chandra, A., Crane, S. J., Tung, E. E., Hanson, G. J., North, F., Cha, S. S., & Takahashi, P. Y. (2015). Patient-reported geriatric symptoms as risk factors for hospitalization and emergency department visits. *Aging and Disease*, 6(3), 188-195. <https://doi.org/10.14336%2FAD.2014.0706>
- Cochrane. (2022). *Data extraction forms*. Cochrane: Developmental, Psychosocial and Learning Problems. <https://dplp.cochrane.org/data-extraction-forms>
- Cochrane Effective Practice and Organisation of Care (EPOC). (2017). *Suggested risk of bias criteria for EPOC reviews: EPOC Resources for review authors*. 2017. https://epoc.cochrane.org/sites/epoc.cochrane.org/files/public/uploads/Resources-for-authors2017/suggested_risk_of_bias_criteria_for_epoc_reviews.pdf
- Davey, P., Brown, E., Charani, E., Fenelon, L., Gould, I. M., Holmes, A., Ramsay, C. R., Wiffen, P. J., & Wilcox, M. (2013). Interventions to improve antibiotic prescribing practices for hospital inpatients. *Cochrane Database of Systematic Reviews*, 2(2), CD003543. <https://doi.org/10.1002/14651858.CD003543.pub3>
- de Gelder, J., Lucke, J. A., de Groot, B., Fogteloo, A. J., Anten, S., Heringhaus, C., Dekkers, O. M., Blauw, G. J., & Mooijaart, S. P. (2018). Predictors and outcomes of revisits in older adults discharged from the emergency department. *Journal of the American Geriatrics Society*, 66(4), 735-741. <https://doi.org/10.1111/jgs.15301>
- Deschodt, M., Devriendt, E., Sabbe, M., Knockaert, D., Deboutte, P., Boonen, S., Flamaing, J., & Milisen, K. (2015). Characteristics of older adults admitted to the emergency department (ED) and their risk factors for ED readmission based on comprehensive geriatric assessment: A prospective cohort study. *BMC Geriatrics*, 15(1), 1-10. <https://doi.org/10.1186/s12877-015-0055-7>
- Dufour, I., Chouinard, M.-C., Dubuc, N., Beaudin, J., Lafontaine, S., & Hudon, C. (2019). Factors associated with frequent use of emergency-department services in a geriatric population: A systematic review. *BMC Geriatrics*, 19(1), 1-9. <https://doi.org/10.1186/s12877-019-1197-9>
- Enguidanos, S., Gibbs, N., & Jamison, P. (2012). From hospital to home: A brief nurse practitioner intervention for vulnerable older adults. *Journal of Gerontological Nursing*, 38(3), 40-50. <https://doi.org/10.3928/00989134-20120116-01>
- Feliciano, E., Feliciano, A., Palompon, D., & Boshra, A. (2022). Aging-related Resiliency Theory Development. *Belitung Nursing Journal*, 8(1), 4-10. <https://doi.org/10.33546/bnj.1631>
- Fougère, B., Aubertin-Leheudre, M., Vellas, B., Andrieu, S., Demougeot, L., Cluzan, C., & Cesari, M. (2016). Clinical research for older adults in rural areas: the MINDED study experience. *Age*, 38(2), 1-4. <https://doi.org/10.1007/s11357-016-9892-3>
- Hasegawa, K., Tsugawa, Y., Tsai, C.-L., Brown, D. F. M., & Camargo, C. A. (2014). Frequent utilization of the emergency department for acute exacerbation of chronic obstructive pulmonary disease. *Respiratory Research*, 15(1), 1-13. <https://doi.org/10.1186/1465-9921-15-40>
- Heeren, P., Devriendt, E., Fieuws, S., Wellens, N. I. H., Deschodt, M., Flamaing, J., Sabbe, M., & Milisen, K. (2019). Unplanned readmission prevention by a geriatric emergency network for transitional care (URGENT): A prospective before-after study. *BMC Geriatrics*, 19(1), 1-10. <https://doi.org/10.1186/s12877-019-1233-9>
- Joanna Briggs Institute. (2020). *Critical appraisal tools*. <https://jbi.global/critical-appraisal-tools>
- Kessler, C., Williams, M. C., Moustoukas, J. N., & Pappas, C. (2013). Transitions of care for the geriatric patient in the emergency department. *Clinics in Geriatric Medicine*, 29(1), 49-69. <https://doi.org/10.1016/j.cger.2012.10.005>
- Kim, J., Lin, A., Absher, R., Makhoul, T., & Wells, C. (2021). Comprehensive and collaborative pharmacist transitions of care service for underserved patients with chronic obstructive pulmonary disease. *Chronic Obstructive Pulmonary Diseases: Journal of the COPD Foundation*, 8(1), 152-161. <https://doi.org/10.15326%2Fjcopdf.2019.0175>

- Lee, I. F.-K., Yau, F. N., Yim, S. S.-H., & Lee, D. T.-F. (2018). Evaluating the impact of a home-based rehabilitation service on older people and their caregivers: A matched-control quasi-experimental study. *Clinical Interventions in Aging*, 13, 1727-1737. <https://doi.org/10.2147%2FICIA.S172871>
- Legrain, S., Tubach, F., Bonnet-Zamponi, D., Lemaire, A., Aquino, J. P., Paillaud, E., Taillandier-Herich, E., Thomas, C., Verny, M., & Pasquet, B. (2011). A new multimodal geriatric discharge-planning intervention to prevent emergency visits and rehospitalizations of older adults: The optimization of medication in AGEd multicenter randomized controlled trial. *Journal of the American Geriatrics Society*, 59(11), 2017-2028. <https://doi.org/10.1111/j.1532-5415.2011.03628.x>
- Liang, H. Y., Hann Lin, L., Yu Chang, C., Mei Wu, F., & Yu, S. (2021). Effectiveness of a nurse-led tele-homecare program for patients with multiple chronic illnesses and a high risk for readmission: A randomized controlled trial. *Journal of Nursing Scholarship*, 53(2), 161-170. <https://doi.org/10.1111/jnu.12622>
- Low, L. L., Vasanwala, F. F., Ng, L. B., Chen, C., Lee, K. H., & Tan, S. Y. (2015). Effectiveness of a transitional home care program in reducing acute hospital utilization: A quasi-experimental study. *BMC Health Services Research*, 15(1), 1-8. <https://doi.org/10.1186/s12913-015-0750-2>
- Lowthian, J., Straney, L. D., Brand, C. A., Barker, A. L., Smit, P. D. V., Newnham, H., Hunter, P., Smith, C., & Cameron, P. A. (2016). Unplanned early return to the emergency department by older patients: The Safe Elderly Emergency Department Discharge (SEED) project. *Age and Ageing*, 45(2), 255-261. <https://doi.org/10.1093/ageing/afv198>
- McPake, B., & Mahal, A. (2017). Addressing the needs of an aging population in the health system: The Australian case. *Health Systems & Reform*, 3(3), 236-247. <https://doi.org/10.1080/23288604.2017.1358796>
- Mi, R., Hollander, M. M., Jones, C., DuGoff, E. H., Caprio, T. V., Cushman, J. T., Kind, A. J. H., Lohmeier, M., & Shah, M. N. (2018). A randomized controlled trial testing the effectiveness of a paramedic-delivered care transitions intervention to reduce emergency department revisits. *BMC Geriatrics*, 18(1), 1-9. <https://doi.org/10.1186/s12877-018-0792-5>
- Naseer, M., Agerholm, J., Fastbom, J., Schön, P., Ehrenberg, A., & Dahlberg, L. (2020). Factors associated with emergency department revisits among older adults in two Swedish regions: A prospective cohort study. *Archives of Gerontology and Geriatrics*, 86, 103960. <https://doi.org/10.1016/j.archger.2019.103960>
- Naylor, M., & Keating, S. A. (2008). Transitional care. *AJN The American Journal of Nursing*, 108(9), 58-63. <https://doi.org/10.1097/01.NAJ.0000336420.34946.3a>
- Naylor, M. D., Aiken, L. H., Kurtzman, E. T., Olds, D. M., & Hirschman, K. B. (2011). The importance of transitional care in achieving health reform. *Health Affairs*, 30(4), 746-754. <https://doi.org/10.1377/hlthaff.2011.0041>
- Occelli, P., Touzet, S., Rabilloud, M., Ganne, C., Poupon Bourdy, S., Galamand, B., Debray, M., Dartiguepeyrou, A., Chuzeville, M., & Comte, B. (2016). Impact of a transition nurse program on the prevention of thirty-day hospital readmissions of elderly patients discharged from short-stay units: Study protocol of the PROUST stepped-wedge cluster randomised trial. *BMC Geriatrics*, 16(1), 1-9. <https://doi.org/10.1186/s12877-016-0233-2>
- Paksopis, T., Suteparuk, C., Sinsuwan, N., Rojsaengroeng, R., Sri-on, J., & Vanichkulbodee, A. (2019). Emergency department visited for elderly patients in Bangkok, Thailand before and after severe flooding in 2011 *Thai Journal of Emergency Medicine*, 1(1), 5-15.
- Pham, H. H., O'Malley, A. S., Bach, P. B., Saiontz-Martinez, C., & Schrag, D. (2009). Primary care physicians' links to other physicians through Medicare patients: The scope of care coordination. *Annals of Internal Medicine*, 150(4), 236-242. <https://doi.org/10.7326/0003-4819-150-4-200902170-00004>
- Rasmussen, L. F., Grode, L. B., Lange, J., Barat, I., & Gregersen, M. (2021). Impact of transitional care interventions on hospital readmissions in older medical patients: A systematic review. *BMJ Open*, 11(1), e040057. <http://dx.doi.org/10.1136/bmjopen-2020-040057>
- Robinson, T. E., Boyd, M. L., North, D., Wignall, J., Dawe, M., McQueen, J., Frey, R. A., Raphael, D. L., & Kerse, N. (2021). Proactive primary care model for frail older people in New Zealand delays aged-residential care: A quasi-experiment. *Journal of the American Geriatrics Society*, 69(6), 1617-1626. <https://doi.org/10.1111/jgs.17064>
- Sabir, F., Tomlinson, J., Strickland-Hodge, B., & Smith, H. (2019). Evaluating the Connect with Pharmacy web-based intervention to reduce hospital readmission for older people. *International Journal of Clinical Pharmacy*, 41(5), 1239-1246. <https://doi.org/10.1007/s11096-019-00887-3>
- Salvi, F., Rossi, L., Lattanzio, F., & Cherubini, A. (2017). Is polypharmacy an independent risk factor for adverse outcomes after an emergency department visit? *Internal and Emergency Medicine*, 12(2), 213-220. <https://doi.org/10.1007/s11739-016-1451-5>
- Santos, F. S. d., Dias, B. M., & Reis, A. M. M. (2019). Emergency department visits of older adults within 30 days of discharge: Analysis from the pharmacotherapy perspective. *Einstein (São Paulo)*, 18. https://doi.org/10.31744/einstein_journal/2020AO4871
- Schapiro, M., Outumuro, M. B., Giber, F., Pino, C., Mattiussi, M., Montero-Odasso, M., Boietti, B., Saimovici, J., Gallo, C., & Hornstein, L. (2021). Geriatric co-management and interdisciplinary transitional care reduced hospital readmissions in frail older patients in Argentina: Results from a randomized controlled trial. *Aging Clinical and Experimental Research*, 34(1), 85-93. <https://doi.org/10.1007/s40520-021-01893-0>
- Schoonover, H., Corbett, C. F., Weeks, D. L., Willson, M. N., & Setter, S. M. (2014). Predicting potential postdischarge adverse drug events and 30-day unplanned hospital readmissions from medication regimen complexity. *Journal of Patient Safety*, 10(4), 186-191. <https://doi.org/10.1097/PTS.0000000000000067>
- Sheikh, S. (2019). Risk factors associated with emergency department recidivism in the older adult. *Western Journal of Emergency Medicine*, 20(6), 931-938. <https://doi.org/10.5811%2Fwestjem.2019.7.43073>
- Skelly, A. C., Dettori, J. R., & Brodt, E. D. (2012). Assessing bias: the importance of considering confounding. *Evidence-Based Spine-Care Journal*, 3(01), 9-12. <https://doi.org/10.1055/s-0031-1298595>
- Son, Y.-J., & You, M.-A. (2015). Transitional care for older adults with chronic illnesses as a vulnerable population: Theoretical framework and future directions in nursing. *Journal of Korean Academy of Nursing*, 45(6), 919-927. <https://doi.org/10.4040/jkan.2015.45.6.919>
- Song, M., Jin, X., Ko, H. N., & Tak, S. H. (2016). Chief complaints of elderly individuals on presentation to emergency department: A retrospective analysis of South Korean National Data 2014. *Asian Nursing Research*, 10(4), 312-317. <https://doi.org/10.1016/j.anr.2016.10.001>
- Ukkonen, M., Jämsen, E., Zeitlin, R., & Pauniahio, S.-L. (2019). Emergency department visits in older patients: A population-based survey. *BMC Emergency Medicine*, 19(1), 1-8. <https://doi.org/10.1186/s12873-019-0236-3>
- Verhaegh, K. J., MacNeil-Vroomen, J. L., Eslami, S., Geerlings, S. E., de Rooij, S. E., & Buurman, B. M. (2014). Transitional care interventions prevent hospital readmissions for adults with chronic illnesses. *Health Affairs*, 33(9), 1531-1539. <https://doi.org/10.1377/hlthaff.2014.0160>
- Wee, S. L., Loke, C. K., Liang, C., Ganesan, G., Wong, L. M., & Cheah, J. (2014). Effectiveness of a national transitional care program in reducing acute care use. *Journal of the American Geriatrics Society*, 62(4), 747-753. <https://doi.org/10.1111/jgs.12750>

Cite this article as: Jehloh, L., Songwathana, P., & Sae-Sia, W. (2022). Transitional care interventions to reduce emergency department visits in older adults: A systematic review. *Belitung Nursing Journal*, 8(3), 187-196. <https://doi.org/10.33546/bnj.2100>