



Belitung Nursing Journal

E-ISSN: 2477-4073 | P-ISSN: 2528-181X

Volume 8, Issue 4
July - August 2022

Edited by Assoc. Prof. Dr. Yupin Aungsuroch & Dr. Joko Gunawan

The Official Publication of
Belitung Raya Foundation
Department of Publication, Indonesia





Examining the impact of upright and recumbent positions on labor outcomes in Saudi Arabia: A quasi-experiment

Zahra Al Aryani¹, Abeer Orabi^{2,3} , and Howieda Fouly^{2,4*} 

¹ NICU Department, King Fahad Hospital Al-Baha, Kingdom of Saudi Arabia

² College of Nursing - Jeddah, King Saud bin Abdulaziz University for Health Sciences, Jeddah, Kingdom of Saudi Arabia

³ Maternal and Newborn Health Nursing Department, Faculty of Nursing, Cairo University, Egypt

⁴ Obstetrics and Gynecology Nursing Department, Assiut University, Egypt

Abstract

Background: Helping the woman to adopt a comfortable position during childbirth significantly affects labor changes and pain management. However, there is a lack of consensus on the impacts of different childbirth positions on labor outcomes. In addition, a scarce of studies have assessed the effects of the upright and recumbent positions on delivery outcomes, especially in the Kingdom of Saudi Arabia.

Objective: This study aimed to examine and compare the influence of upright and recumbent childbirth positions on birth outcomes.

Methods: The setting was the childbirth unit at East Jeddah Hospital from November 2020 to March 2021. The research design was quasi-experimental, including 300 women in labor under 18-45 years. The sample includes two equal groups of 150 women: upright (experiment) and recumbent (comparison) position groups. Three tools were used to collect data: a structured interviewing questionnaire, the modified WHO partograph, and the Wong-Baker FACES® pain rating scale. Descriptive statistics, chi-square, independent *t*-test, and paired *t*-test were used for data analysis.

Results: Women in the recumbent position spent a longer duration in the first, second, and third stages of childbirth and had higher pain scores and less satisfaction with the assumed position than women in the upright position, with a highly significant difference ($p < 0.001$).

Conclusion: Laboring women in upright positions experienced faster progress of labor, shorter duration of childbirth, less pain, and higher satisfaction than those assumed recumbent positions. This study serves as an input for midwives and nurses to enable them to offer appropriate advice to improve intrapartum care. In addition, educational programs targeting pregnant women about the positive impacts of upright position on women's birth experience are encouraged.

Keywords

childbirth; positions; labor outcomes; nursing; midwifery; Saudi Arabia

*Corresponding author:

Associate professor. Dr. Howieda Fouly
College of Nursing-KSAU-HS. Jeddah KSA
Faculty of Nursing- Assiut University, Egypt
Emails: hoida.elfouly2@aun.edu.eg |
hoida_elfouly@yahoo.com

Article info:

Received: 4 April 2022

Revised: 6 May 2022

Accepted: 18 July 2022



This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License, which allows others to remix, tweak, and build upon the work non-commercially as long as the original work is properly cited. The new creations are not necessarily licensed under the identical terms.

E-ISSN: 2477-4073 | P-ISSN: 2528-181X

Background

Childbirth is a physiological process that involves a sequential, gradual, and integrated set of changes within the birth canal to end with the delivery of the fetus and the placenta (Modoor et al., 2021; Ricci, 2012). Putting laboring women in a relaxed position is vital to help them to deal well with these changes, reduce pain and facilitate normal delivery (White et al., 2011). Moreover, maternal positioning during childbirth affects childbirth progress. At the same time, it can reduce the need for the induction or augmentation of delivery and manage pain and stress (Watson & Cooke, 2018).

Positions for women in childbirth can be categorized into upright positions as standing, sitting, kneeling, squatting, walking, and on all fours, and recumbent positions as lateral, supine, and semi-recumbent (Kemp et al., 2013). In the upright

position, gravity reduces aortocaval compression, which makes the uterine contractions more effective and ensures better fetal alignment in the birth canal (Gizzo et al., 2014). In this context, the results from an experimental study demonstrated that adopting an upright position in the first stage of childbirth, when the cervix was dilated 2–4 cm, reduced pain, improved the effectiveness of uterine contractions, decreased the rate of cesarean birth, perineal trauma, an episiotomy, and significantly decreased the use of analgesics (Deliktas & Kukulu, 2018). Furthermore, Emam and Al-Zahrani (2018) reported that using the upright position positively impacted the progress of childbirth by increasing the effectiveness of the uterine contractions and facilitating cervical dilatation and fetal descent, compared to the recumbent position, and this finding had statistical significance. In contrast, the adoption of the upright position throughout childbirth was not shown to have a statistically

significant impact on the duration and mode of delivery (Deliktas & Kukul, 2018).

Elsewhere, the recumbent position may be used by women during childbirth because of being recommended at admission (Mselle & Eustace, 2020). Using a recumbent position during birth may be seen as beneficial by some healthcare providers to palpate the maternal abdomen, assess uterine contractions, presentation, and position, listen to the fetal heart rate (FHR), and conduct vaginal exams efficiently. However, it can inhibit the progress of childbirth as the gravid uterus compresses the abdominal blood vessels and impairs circulation, including uteroplacental perfusion, which may adversely affect maternal and fetal well-being (Cunningham et al., 2009).

More and more people worldwide are paying attention to the standard of maternity care. Further, it is evaluated against using updated and evidence-based hospital guidelines for birth management (Altaweli et al., 2014). The selection of a maternal birth position is a vital aspect of midwifery care during childbirth; however, in many cases, it may be neglected by some healthcare providers. Adopting a recumbent position in delivery is a common practice in the current century, although it should only be adopted when indicated (Desseauve et al., 2017).

There is a lack of consensus on the impacts of different maternal positions during childbirth on maternal, fetal, and neonatal outcomes. Available evidence in this field is controversial and fragmented. In addition, a scarce of studies have assessed the effects of the upright and recumbent positions on childbirth outcomes during the first stage of childbirth, especially in the Kingdom of Saudi Arabia (Emam & Al-Zahrani, 2018). Thus, it is essential to identify other maternal birth positions that could be used during childbirth's first stage and understand how they impact childbirth outcomes.

Women giving birth must comprehend the advantages and disadvantages of each position and be encouraged to choose a position that will provide them with the most incredible comfort, control, and benefits (Lawrence et al., 2013). In this regard, midwives play an important role in collaborating with other healthcare professionals regarding providing adequate care to women in childbirth. To help laboring women make informed decisions regarding their positions, nurses or midwives must provide concise, consistent, and evidence-based descriptions of the hazards and advantages of each position (Cunningham et al., 2009; Oats & Abraham, 2017). In addition, nurses and midwives should consider each pregnant woman's feelings and identify the obstetric factors that may interfere with the chosen birthing position (Priddis et al., 2012). One of the critical areas of practice is the mother's position during the early labor stage. Despite an increasing body of evidence demonstrating the physical advantages for women and fetuses when they use an upright posture during childbirth, the majority of women around the globe give birth to their children while laying on their backs in bed, with a few notable exceptions, a practice that is not supported by empirical evidence.

In the past, the use of the supine position was encouraged by traditional and former practitioners. Since then, the benefits of the upright position have been revealed, and research backs them up; this position has subsequently been encouraged by some midwives and obstetricians, and it is

requested by women in childbirth (Gizzo et al., 2014). Unfortunately, the impact of upright versus recumbent postures during the first stage of birth on the mother, fetus, and newborn is rarely agreed upon, and the evidence in this field is frequently disputed and fragmentary.

The World Health Organization (WHO) ended with no evidence to back up using a recumbent posture in early labor. However, it is shown that upright positions during early labor reduce the need for interference and the duration of childbirth while having little impact on the well-being of the mother and fetus. Consequently, midwives and nurses should allow women to assume whichever position is most comfortable for them during the first stage of childbirth (World Health Organization, 2018).

Considering that few studies had addressed the impact of different postures on childbirth outcomes in Saudi Arabia, the current research aimed to evaluate the effect of the upright compared to the recumbent postures on birth outcomes in Jeddah. There are four hypotheses for this study; it was assumed that women who adopted the upright position during first-stage childbirth would experience (1) faster progress with their childbirth, (2) shorter childbirth duration, (3) less pain, and (4) more satisfaction with this position than women who used the recumbent posture.

Methods

Study Design

The study used a quasi-experimental design.

Participants/Samples

The setting for this study was the childbirth unit at East Jeddah Hospital, under the auspices of the Ministry of Health of KSA.

The inclusion criteria of the sample were women in the active phase of first-stage childbirth, aged 18–45 years, with normal body mass index, pregnant with a single fetus with the cephalic presentation, term pregnancy (38–41 weeks), and had the willingness to be part of the study. While high-risk pregnant women (Gestational diabetic Mellitus, cardiac disease, pre-eclampsia placenta previa & abruption placenta), elective cesarean section, malposition, induced or augmented childbirth, meconium-stained amniotic fluid, and epidural anesthesia were excluded.

The sample size was estimated to be 300 using an online Raosoft® sample calculator (<http://www.raosoft.com>). The input data were a population size of 1,267 over six months (Ministry of Health of Saudi Arabia, 2019), 5.0% acceptable error of margin, 95% level of confidence, and 50% response rate.

Purposive sampling was used to recruit 300 eligible laboring women. Afterward, the participants were categorized into two groups, 150 each: upright position (Study group) and recumbent position as routinely used for maternity care in the study setting (Comparison group). Twenty-five women were excluded after recruitment owing to the presence of meconium-stained amniotic fluid, the administration of pethidine, withdrawal from the study, or language barriers (Figure 1).




Figure 1 Flow chart of the sampling procedure used when recruiting the study's participants

Instruments

A structured questionnaire: the modified WHO partograph and the Wong-Baker FACES® Pain Rating Scale were used to obtain the needed data. The first part of the questionnaire elicited information on the sociodemographic background of the participants, such as age, residence, occupation, educational level, gestational age, gravidity, and parity.

The second part assessed maternal satisfaction with the adopted position. If a mother would use it in the subsequent childbirth through two questions, their answers were evaluated by a 5-points Likert scale from 0–4 (Emam & Al-Zahrani, 2018). Permission to use this part was obtained from the corresponding author by email.

The Modified World Health Organization partograph was used to graphically determine childbirth progress using data pertaining to dilatation of the cervix, fetal descent, and contractions of the uterus (frequency, duration & intensity), the

duration of the first, second, and third stages of childbirth and the method of delivery (World Health Organization, 2018).

The Wong-Baker FACES Foundation (2019) was used after the researchers explained the scoring system to the participants. Then, they were requested to select the best face that depicted the experienced pain.

The validity of the data collection tools was ensured by a review of experts in the field of obstetrics and gynecology who assessed their relevancy and appropriateness to meet the study's aim. Reliability was evaluated by pilot testing of the instruments to ensure their consistency. A pilot study was conducted by pre-testing the tools on 10% of the sample (30 women in childbirth) eligible for the study to assess their applicability, consistency, and accuracy and to determine the needed time to complete the research and potential challenges that may hinder the data collection (Polit & Beck, 2018).

The pilot study data were reviewed and subjected to statistical analysis. The total reliability of the data measurement scale was 0.63.

Data Collection

Data collection was completed in five months (November 2020–March 2021). Data collection started by explaining the aim and objectives of the study to nurses/midwives in the childbirth unit at East Jeddah Hospital. Then, the researchers selected two rooms for the comparison group and two for the experiment group. Next, data were collected through interviews, assessments, implementation, and evaluation phases (Figure 2).




Figure 2 The data collection process

Interview phase. The researcher explained the aim and study details to women before obtaining their consent. After that, the participants were interviewed to collect sociodemographic data.

Assessment phase. The standard data on childbirth progress, such as contractions of the uterus (frequency, duration & strength), dilatation of the cervix, fetal descent, and pain levels, were collected upon the admission of the participants to the unit.

Implementation phase. While in childbirth, the participants in both groups received intravenous fluids, back massage, and prescribed paracetamol. In addition to testing for COVID-19 as routine care in the first stage of childbirth, according to the hospital policy. The researchers explained the different types of an upright position and their benefits to each woman in the study group; then, the participant was requested to adopt an upright position of choice for 15–20 minutes each hour and return to bed in between for 10–15 minutes. The selected positions included sitting on a birthing ball with support, on all

fours, and squatting. Participants in the comparison group adopted either a supine or lateral position.

Evaluation phase. The researchers evaluated the effects of the upright and recumbent positions on childbirth progress hourly using the data collection tools in terms of the interval, frequency, and duration of contractions of the uterus, dilatation of the cervix, and fetal descent, as well as pain levels and maternal satisfaction with the adopted position.

Data Analysis

The Statistical Package managed data for Social Sciences®, version 25 for Windows®. Frequencies, percentages, means, and standard deviations were used for data presentation. Chi-square, independent *t*-test, and paired *t*-test were also used to assess the association and significance of the finding between the study variables and to test the study hypotheses. A significant difference was set at ≤ 0.05 .

Ethical Considerations

The research was performed after being reviewed and approved by the Research Unit in the College of Nursing, the King Abdullah International Medical Research Center, and the institutional review board (IRB) at King Abdulaziz Medical City, Western Region, Jeddah, KSA. Approval from the IRB of the

Ministry of Health, KSA, was also granted, and the administrative personnel in East Jeddah Hospital were contacted before beginning data collection. The researcher clarified the study's aim and nature to women who met the criteria for inclusion and obtained their informed consent for participation. It was clear to the women that their involvement in the study was voluntary, and they had the right to revoke it at any moment. Privacy and anonymity were kept.

Results

The mean age of the participants in the experiment group was 32.20 ± 28.37 years, with 43.3% of them aged 26 to 30 years as compared to 30.50 ± 5.3 years, with 40.0% of them aged 31 to 35 years in the comparison group. In addition, in the experiment group, 74.7% received university or higher education, 62% were housewives, 54.7% were living in urban areas, and 92.7% were Saudi, compared to 62.7%, 55.3%, 44.7%, and 90.7%, respectively in the comparison group. Finally, the mean gestational age during delivery was 39.1 ± 2.93 weeks in the experiment group compared to 39.2 ± 3.99 weeks in the comparison group which reflected no statistically significant differences between the two groups (Table 1).

Table 1 Distribution of the sociodemographic data among the experiment and comparison groups ($n = 150$ for each group)

Variable	Upright position (experiment group)		Recumbent position (comparison group)	
	<i>n</i>	%	<i>n</i>	%
Age in years				
20 to 25	31	20.7	39	26.0
26 to 30	65	43.3	28	18.7
31 to 35	43	28.7	60	40.0
≥35	11	7.3	23	15.3
Mean ± SD	32.20 ± 28.37		30.50 ± 5.3	
x²= 377.43 <i>p</i> = 0.49				
Educational level				
Secondary	38	25.3	56	37.3
University or higher	112	74.7	94	62.7
x²= 1.19 <i>p</i> = 0.27				
Occupation				
Work-related to the health sector	14	9.3	9	6.0
Work not related to the health sector	43	28.7	58	38.7
Housewife	93	62.0	83	55.3
x²= 3.09 <i>p</i> = 0.54				
Residence				
Rural	68	45.3	83	55.3
Urban	82	54.7	67	44.7
x²= 1.04 <i>p</i> = 0.23				
Nationality				
Saudi	139	92.7	136	90.7
Non-Saudi	11	7.3	14	9.3
x²= 0.001 <i>p</i> = 0.97				
Gestational age in weeks				
Mean ± SD	39.1 ± 2.93		39.1 ± 2.93	
x²= 544.11 <i>p</i> = 0.94				

The mean baseline duration of uterine contractions in seconds among the participants in the experiment group was 36.76 ± 7.9 compared to 38.76 ± 10.63 in the comparison group and reflected no statistically significant difference ($p < 0.06$).

After the interventions, the mean duration of uterine contractions was higher among participants in the experimental group than among the participants in the recumbent position, reflecting a statistically significant difference in childbirth outcomes between the two groups, particularly after two hours ($t_{298} = 92.10$, $p = 0.001$), three

hours ($t_{248} = 240.83$, $p = 0.001$), and four hours ($t_{12} = 2.11$, $p = 0.04$).

In addition, the intervals between uterine contractions reflect a statistically significant difference after two hours ($t_{298} = 2.68$, $p = 0.001$), three hours ($t_{248} = 2.23$, $p = 0.03$), and four hours ($t_{280} = 12$, $p = 0.02$), respectively. Also, there was a statistically significant difference in the number of uterine contractions per 10 minutes, especially after one hour ($t_{298} =$

3.07, $p = 0.001$), two hours ($t_{298} = 3.69$, $p = 0.001$), and three hours ($t_{248} = 4.12$, $p = 0.001$), but not after four hours ($t_{12} = 1.55$, $p = 0.16$) (Table 2).

In addition, Table 3 shows that there was a significant difference in the intensity of uterine contraction between the experimental and control groups, especially after two hours ($X^2 = 5.03$, $p = 0.02$) and three hours ($X^2 = 23.08$, $p = 0.001$).

Table 2 Comparison between the different positions among participants based on the characteristics of their uterine contraction ($n = 150$ for each group)

Variable	Upright position (experiment)		Recumbent position (Comparison)		Independent t- test	df	p-value
	n	Mean ± SD	n	Mean ± SD			
Duration of uterine contractions in seconds							
Baseline	150	36.76 ± 7.9	150	38.76 ± 10.63	1.83	298	0.07
After one hour	150	44.46 ± 9.8	150	42.00 ± 12.28	1.84	298	0.67
After two hours	150	58.00 ± 6.02	150	51.80 ± 11.81	92.10	298	0.001*
After three hours	125	60.00 ± 0.00	125	57.69 ± 5.77	240.83	248	0.001*
After four hours	7	60.00 ± 5.90	7	60.89 ± 5.99	2.11	12	0.04*
Intervals between uterine contractions in minutes							
Baseline	150	3.50 ± 0.63	150	3.54 ± 0.63	0.56	298	0.57
After one hour	150	3.18 ± 0.50	150	3.28 ± 0.50	1.77	298	0.07
After two hours	150	2.97 ± 0.25	150	3.09 ± 0.49	2.68	298	0.001*
After three hours	125	2.76 ± 0.33	125	2.86 ± 0.34	2.23	248	0.03*
After four hours	7	2.5 ± 0.04	7	2.85 ± 0.23	2.80	12	0.02*
Number of uterine contractions per 10 minutes							
Baseline	150	2.60 ± 0.49	150	2.56 ± 0.49	1.08	298	0.278
After one hour	150	2.92 ± 0.54	150	2.74 ± 0.436	3.07	298	0.001*
After two hours	150	3.18 ± 0.39	150	3.01 ± 0.46	3.69	298	0.001*
After three hours	125	3.56 ± 0.49	125	3.28 ± 0.53	4.12	248	0.001*
After four hours	7	3.14 ± 0.37	7	3.29 ± 0.46	1.55	12	0.16

*Statistically significant difference ($p < 0.05$)

Table 3 Comparison between the different groups based on the intensity of uterine contraction ($n = 150$ for each group)

The intensity of uterine contraction		Upright position (experiment)		Recumbent position (comparison)		Test of significant	p-value
		n	%	n	%		
Baseline	Mild	6	4.0	1	0.7	$X^2 = 1.99$	0.13
	Moderate	115	76.7	115	76.7		
	Severe	29	19.3	34	22.7		
After 1 hour	Moderate	88	58.7	73	48.7	$X^2 = 3.01$	0.08
	Severe	62	41.3	77	51.3		
After 2 hours	Moderate	16	10.7	30	20.0	$X^2 = 5.03$	0.02*
	Severe	134	89.3	120	80.0		
After 3 hours	Moderate	1	0.7	2	1.3	$X^2 = 23.08$	0.001*
	Severe	124	99.3	123	82.0		
After 4 hours	Moderate	0	0	1	16.66	$X^2 = 0.74$	0.38
	Severe	7	100	6	83.34		

* Statistically significant difference ($p < 0.05$)

Table 4 shows that there was greater progress in cervical dilatation after three hours among the participants in the upright position group than among those in the recumbent position group, with a highly statistically significant difference between the two groups ($t_{248} = 2.6$, $p < 0.001$). Similarly, there was progress of foetal descent was greater among the participants in the upright position group than among those in the recumbent position group, with a high statistically

significant difference especially after one hour ($t_{298} = 7.04$, $p < 0.001$), two hours ($t_{298} = 9.55$, $p < 0.001$), three hours ($t_{248} = 13.97$, $p < 0.001$), and four hours ($t_{12} = 0.00$, $p < 0.02$), respectively. In addition, there was a significant difference in pain scores between the two groups after one hour ($t_{298} = 18.52$, $p < 0.001$), two hours ($t_{298} = 90.81$, $p < 0.001$), and three hours ($t_{248} = 109.38$, $p < 0.001$), respectively (Table 4).

Table 4 Comparison between the different positions in relation to cervical dilatation (cm), the descent of the fetal head, and mean pain scores ($n = 150$ for each group)

Variable	Upright position (experiment)		Recumbent position (comparison)		Independent t-test	df	p-value
	n	Mean ± SD	n	Mean ± SD			
Cervical dilatation (cm)							
Baseline	150	5.31 ± 0.80	150	5.30 ± 0.85	0.73	298	0.44
After one hour	150	6.28 ± 0.58	150	6.36 ± 0.92	1.01	298	0.32
After two hours	150	7.9 ± 0.72	150	7.84 ± 1.11	0.57	298	0.54
After three hours	125	9.6 ± 0.69	125	9.36 ± 1.02	2.6	248	0.001*
After four hours	7	10 ± 0.00	7	9.53 ± 0.71	1.01	12	0.34
Fetal head descent/fifths							
Baseline	150	-2.52 ± 0.69	150	-2.42 ± 0.61	1.18	298	0.22
After one hour	150	-1.66 ± 0.68	150	-2.21 ± 0.71	7.04	298	0.001*
After two hours	150	-0.35 ± 0.60	150	-1.21 ± 0.92	9.55	298	0.001 *
After three hours	125	1.18 ± 0.87	125	0.32 ± 0.81	13.97	248	0.001 *
After four hours	7	0.33 ± 0.57	7	0.33 ± 0.57	0.00	12	0.02 *
Mean pain scores							
Baseline	150	3.05 ± 1.00	150	3.09 ± 0.99	0.55	298	0.55
After one hour	150	5.02 ± 1.04	150	6.69 ± 1.11	18.52	298	0.001*
After two hours	150	6.7 ± 1.13	150	8.61 ± 1.08	90.81	298	0.001 *
After three hours	125	8.13 ± 1.24	125	9.08 ± 1.00	109.38	248	0.001*
After four hours	7	8.38 ± 1.24	7	9.00 ± 1.00	108.99	12	0.51

*Statistically significant difference ($p < 0.05$)

Table 5 shows that the duration of the first stage was under 5-6 hours for 74.64% of the experiment group compared to 61.34% of the comparison group. Only 0.66% of the experiment group had >7 hours in the first stage of childbirth compared to 24.66% of the comparison group. The mean duration of the first stage of birth in hours among participants in the experiment and comparison groups was 5.88 ± 0.29 and 6.27 ± 0.68 , respectively, with a highly statistically significant

difference at ($t_{202} = 6.22$, $p = 0.001$). The second stage of childbirth duration was <30 minutes for 80% of participants in the experiment group compared to 76% in the comparison group. There was a statistically significant difference between the two groups regarding the duration of the second stage of childbirth ($t_{32} = 3.44$, $p = 0.001$) and the third stage of birth ($t_{72} = 14.92$, $p = 0.001$).

Table 5 Comparison between the different positions based on the duration of childbirth ($n = 150$ for each group)

Variable	Upright position		Recumbent position		Independent t-test	df	p-value
	n	%	n	%			
Duration of the first stage of childbirth in hours							
5-6	112	74.6	92	61.3	6.22	202	0.001 **
6-7	37	24.7	21	14.00		56	
>7	1	0.6	37	24.6		36	
Mean ± SD	5.88 ± 0.29		6.27 ± 0.68				
Duration of the second stage of childbirth in minutes							
<30	120	80.0	114	76.0	3.44	232	0.001**
30-60	30	20.0	33	22.0		61	
>60	0	0.0	3	2.0		1	
Mean ± SD	25.3 ± 9.11		29.30 ± 10.33				
Duration of the third stage of childbirth in minutes							
10-20	145	96.7	129	86.0	14.92	272	0.001 **
20-30	5	3.3	21	14.0		24	
Mean ± SD	10.33 ± 1.80		15.83 ± 4.07				

Table 6 shows statistically significant differences between the two groups with the type of membrane rupture ($X^2 = 27.4$, $p = 0.001$). Of the participants in the experiment group, 57.3% had normal vaginal delivery compared to 44.70% of participants in the comparison group, with a highly statistically significant difference at ($X^2 = 7.42$, $p = 0.001$).

In addition, **Table 7** shows that the mean score satisfaction with the assumed position was higher among the experimental

group (4.03 ± 0.365) than among the comparison group (2.63 ± 1.303), reflecting a statistically significant difference ($t_{298} = 13.01$, $p = 0.001$). Additionally, the mean score for preference of the assumed position in the subsequent childbirth was higher in the experimental group (4.32 ± 0.771) than in the comparison group (2.63 ± 0.628), with a statistically significant difference ($t_{298} = 21.77$, $p = 0.001$).

Table 6 Comparison between the different positions based on the type of membrane rupture and mode of delivery ($n = 150$ for each group)

Variable	Upright position		Recumbent position		Test of significant	p-value
	n	%	n	%		
Type of membranes rupture						
AROM	79	52.7	72	48.0	X ² = 27.4	0.001*
SROM	71	47.3	78	52.0		
Type of vaginal delivery						
Normal	86	57.3	67	44.7	X ² = 7.42	0.001*
With episiotomy	0	0.0	3	1.0		
With 1st degree of perineal tear	49	32.7	32	21.3		
With 2nd degree of perineal tear	15	10.0	47	32.7		
Ventose delivery	0	0.0	1	0.3		

Table 7 Comparison between the upright and recumbent position groups based on maternal satisfaction with the assumed position

Maternal satisfaction with the assumed position	Experiment group		Comparison group		Independent t-test	df	p-value
	n	Mean \pm SD	n	Mean \pm SD			
Level of satisfaction	150	4.03 \pm 0.365	150	2.63 \pm 1.303	13.01	298	0.001 *
Position preference in the next childbirth	150	4.32 \pm 0.771	150	2.63 \pm 0.628	21.77	298	0.001 *

*Statistically significant difference ($p < 0.01$)

Discussion

Findings revealed that participants in both groups were demographically homogeneous, with no statistically significant differences. These results correspond with the outcomes of an observational cohort study conducted by Gizzo et al. (2014) and a quasi-experimental study by Emam and Al-Zahrani (2018) to compare the impacts of recumbent and alternative positions on childbirth progress, mode of delivery, and neonatal well-being. In addition, the findings revealed no substantial differences between both groups in relation to age, educational level, nationality, residence, and occupation. This uniformity among the experiment groups helps reduce extraneous factors, which may interrupt the accuracy of measuring the effects of the planned intervention on the childbirth process and outcome.

We discuss the findings based on the labor outcomes: childbirth progress, duration of childbirth, mode of delivery, maternal satisfaction, and level of pain.

Childbirth progress. The mean duration of uterine contractions was higher among participants in the experimental group than among the participants in the comparison group, with statistically significant differences between the participant groups. Similarly, in an Egyptian study, Emam and Al-Zahrani (2018) reported an upright position's positive effect on childbirth progress. These might be related to the effect of gravity in an upright position on preventing aortocaval squeezing and thereby strengthening uterine contractions. Also, the findings of our study revealed a high statistically significant difference in cervical dilatation between the two groups. Likewise, previous studies by Gizzo et al. (2014) and Chaillet et al. (2014) found that women in the upright position group showed better progress in cervical dilatation than those in the recumbent position group after the first, second, and third hours of assuming the position. These results may be attributed to the fact that upright positions such as sitting, standing, and kneeling during the first stage of childbirth enable the abdominal wall to relax, thereby enabling

gravity to pull the uterine fundus forward. In addition, concerning the fetal head descent, the present study found a significant difference between the two groups at the first, second, and third hours of childbirth after assuming the position. Similarly, Gizzo et al. (2014) and Emam and Al-Zahrani (2018) revealed that upright positions aid in bringing the fetus down into the birth canal with the help of gravity. That is probably because upright positions strengthen the pelvic muscles and expand the pelvic area, which improves fetal descent.

Duration of childbirth. The current study's findings align with Berta et al. (2019) study, which investigated the effect of maternal birth positions on the duration of the second stage of childbirth by conducting a systematic review and meta-analysis of databases. They reported a reduction in the duration of the second stage of birth among women in a relaxed sacrum birthing position. In addition, these findings are consistent with previous studies (Walker et al., 2018; Zaky, 2016), which reported that women in the upright position group had a significantly shorter duration of the first, second, and third stages of childbirth than women in the other group. Furthermore, it was suggested that women who assumed upright had a shorter childbirth duration by 1 hour and 22 minutes than women who assumed the recumbent position. They justified these findings by referring to changing maternal position, frequently moving the pelvic bones, and assisting the fetal descent into the pelvic canal. In contrast, Huang et al. (2019) reported that the upright posture during the first stage of childbirth did not shorten childbirth duration.

Mode of delivery. The current study showed that most participants in the experiment group had normal vaginal delivery compared to a quarter of the participants in the comparison group. Among the participants in the comparison group, one woman had a ventouse delivery because of delayed progress in cervical dilatation and poor maternal effort. Similarly, in two previous studies by Emam and Al-Zahrani (2018) and Deliktas and Kukulu (2018), the majority of women who assumed upright positions in the first stage of childbirth had a normal vaginal delivery, and one-tenth had

forceps and ventouse deliveries in the recumbent position group compared to no one in the upright position group. On the other hand, a study by Ibrahim et al. (2020) found no significant difference between women assigned to upright versus recumbent positions for the mode of delivery and duration of the second and third stages of childbirth.

Maternal satisfaction. The present study finding revealed that participants in the experiment group were more satisfied than those in the comparison group, with a statistically significant difference. The study results are consistent with the findings of studies by Deliktas and Kukulu (2018) and Johnson et al. (2017), who examined the effectiveness of ambulation during the first stage of childbirth on the outcomes of delivery among primigravida women and found that mothers who stayed in bed and did not walk around had lower satisfaction with childbirth than mothers who walked around or moved from one place to another. Similarly, studies by Bohren et al. (2017) and Mathew et al. (2012) reported that women who were allowed to assume an upright posture were more satisfied and secure in childbirth than women who stayed in bed and did not walk around. In the same perspective, a randomized controlled trial conducted by Barasinski et al. (2018) to evaluate the effect of an upright position during the first stage of childbirth on parturients' satisfaction indicated that women in both groups preferred the upright posture. Additionally, Al-Seady et al. (2017) assessed the effect of the upright versus recumbent positions during the active phase of the first stage of childbirth outcomes. Their study revealed that those in the upright position group had higher satisfaction scores than those in the recumbent position group. Moreover, Huang et al. (2019) and Mathew et al. (2012) reported that the upright posture during the first stage of childbirth was a safe and well-accepted alternative for women.

Level of pain. There was a high statistically significant difference between the study and comparison groups in the first, second, third, and fourth hours after assuming the chosen position in relation to pain level. These findings are similar to those of previous studies (Emam & Al-Zahrani, 2018; Shenbagavalli & Menaka, 2016). The justification for this may be that movement can minimize the pain's severity; thus, it was recommended that the woman move freely and respond to her body's signals for mobilization and position modification during childbirth. Moreover, Shenbagavalli and Menaka (2016) revealed that breathing exercises, massage, and position modifications were effective in reducing maternal pain during the first stage of childbirth.

Implications of the Study

With the findings of the current study, some recommendations were developed, including: First, educational programs for midwives and nurses on the different delivery positions, as well as the impacts of each, are essential to enable them to offer appropriate advice to childbirth women and improve intrapartum care. Second, assuming upright positions during the first stage of childbirth must be recommended to all women after explaining the benefits of each one. Third, laboring women must be supported in making an informed choice for their birthing position for a better childbirth experience and satisfaction. Fourth, increasing community awareness by developing orientation and education programs regarding delivery preparation through variable platforms. Fifth, further

research studies and systematic reviews can be conducted with adequate attention to other possible delivery positions and integrating neonatal outcomes. Moreover, assess the nurses' and midwives' knowledge, attitudes, practices, and barriers to applying for different positions during the first stage of childbirth.

Limitations of the Study

The main limitation was the extension period for data collection due to the COVID-19 pandemic in the summer of 2020. However, these limitations did not affect the accuracy of the results.

Conclusion

Based on the findings of the current study, it can be concluded that assuming an upright position during the first stage of childbirth has benefits for a laboring woman by significantly improving childbirth progress, shortening the duration of the three stages of birth, reducing the level of pain, increasing maternal satisfaction with childbirth.

Declaration of Conflicting Interest

The authors declare that there are no conflicting interests.

Funding

The authors received no financial support for this article's research, authorship, and publication.

Acknowledgment

The authors thank the nurses and physicians from East Jeddah Hospital for their outstanding collaboration during data collection.

Authors' Contributions

Conceptualization (AO & HF), Investigation (ZA), Methodology, Project Supervision and Administration (AO & HF), Validation (AO, HF, ZA), Writing – original draft (AO & ZA), Review & Editing (AO & HF). All authors agreed to be accountable in every stage of the study and agreed with the final version of the article to be published.

Authors' Biographies

Zahra Al Aryani, Master Midwifery, is a Head Nurse at the NICU Department, King Fahad Hospital Al-Baha, Kingdom of Saudi Arabia.

Abeer Orabi, PhD, is an Assistant Professor at the College of Nursing - Jeddah, King Saud bin Abdulaziz University for Health Sciences, Jeddah, Kingdom of Saudi Arabia. She is also a Lecturer at the Maternal and Newborn Health Nursing Department, Faculty of Nursing, Cairo University, Egypt.

Howieda Fouly PhD, is an Assistant Professor at the College of Nursing - Jeddah, King Saud bin Abdulaziz University for Health Sciences, Jeddah, Kingdom of Saudi Arabia. She is also an Associate Professor at the Obstetrics and Gynecology Nursing Department, Assiut University, Egypt.

Data Availability

The datasets generated during and analyzed during the current study are available from the corresponding author on reasonable request.

References

- Al-Seady, M. Y., Fadel, E. A., El-Gohary, A. M., & Marzouk, T. (2017). Labour pain and satisfaction of primipara assume upright versus recumbent positions during first stage of labour. *IOSR Journal of Nursing and Health Science* 6(4), 24-30. <https://doi.org/10.9790/1959-0604022430>
- Altawell, R. F., McCourt, C., & Baron, M. (2014). Childbirth care practices in public sector facilities in Jeddah, Saudi Arabia: A descriptive study. *Midwifery*, 30(7), 899-909. <https://doi.org/10.1016/j.midw.2014.03.006>

- Barasinski, C., Debost-Legrand, A., Lemery, D., & Vendittelli, F. (2018). Positions during the first stage and the passive second stage of labor: A survey of French midwives. *Midwifery*, 56, 79-85. <https://doi.org/10.1016/j.midw.2017.10.010>
- Berta, M., Lindgren, H., Christensson, K., Mekonnen, S., & Adefris, M. (2019). Effect of maternal birth positions on duration of second stage of labor: Systematic review and meta-analysis. *BMC Pregnancy and Childbirth*, 19(1), 1-8. <https://doi.org/10.1186/s12884-019-2620-0>
- Bohren, M. A., Hofmeyr, G. J., Sakala, C., Fukuzawa, R. K., & Cuthbert, A. (2017). Continuous support for women during childbirth. *Cochrane Database of Systematic Reviews*(7), CD003766. <https://doi.org/10.1002/14651858.CD003766.pub6>
- Chaillet, N., Belaid, L., Crochetiere, C., Roy, L., Gagné, G. P., Moutquin, J. M., Rossignol, M., Dugas, M., Wassef, M., & Bonapace, J. (2014). Nonpharmacologic approaches for pain management during labor compared with usual care: A meta-analysis. *Birth*, 41(2), 122-137. <https://doi.org/10.1111/birt.12103>
- Cunningham, G., Leveno, K. J., Bloom, S. L., Hauth, J. C., & Rouse, D. (2009). *Williams obstetrics* (23rd ed.). New York: McGraw Hill Professional.
- Deliktas, A., & Kukul, K. (2018). The effect of upright positions during the first stage of labour on childbirth types: A meta-analysis [Report]. *Clinical and Experimental Health Sciences*, 8(2), 128-137. <https://link.gale.com/apps/doc/A558533152/AONE?u=anon~c272f9a&sid=googleScholar&id=2868f0ae>
- Desseauve, D., Fradet, L., Lacouture, P., & Pierre, F. (2017). Position for labor and birth: State of knowledge and biomechanical perspectives. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 208, 46-54. <https://doi.org/10.1016/j.ejogrb.2016.11.006>
- Emam, A. M., & Al-Zahrani, A. E. (2018). Upright versus recumbent position during first stage of labor among primipara women on labor outcomes. *Journal of Nursing Education and Practice*, 8(7), 113-124. <https://doi.org/10.5430/jnep.v8n7p113>
- Gizzo, S., Di Gangi, S., Noventa, M., Bacile, V., Zambon, A., & Nardelli, G. B. (2014). Women's choice of positions during labour: Return to the past or a modern way to give birth? A cohort study in Italy. *BioMed Research International*, 638093. <https://doi.org/10.1155/2014/638093>
- Huang, J., Zang, Y., Ren, L.-H., Li, F.-J., & Lu, H. (2019). A review and comparison of common maternal positions during the second-stage of labor. *International Journal of Nursing Sciences*, 6(4), 460-467. <https://doi.org/10.1016/j.ijnss.2019.06.007>
- Ibrahim, H. A.-F., Said, H. I. I., & Elgzar, W. T. I. (2020). Effect of upright and ambulant positions versus lying down during the active first stage of labor on birth outcomes among nulliparous women: Randomized controlled clinical trial. *Frontiers of Nursing*, 7(3), 239-248. <https://doi.org/10.2478/fon-2020-0035>
- Johnson, N., Nokomis, Z., & Stark, M. A. (2017). The nurses' role in providing comfort during childbirth using ambulation and hydrotherapy. *International Journal of Studies in Nursing*, 3(1), 123-131. <https://doi.org/10.20849/ijns.v3i1.347>
- Kemp, E., Kingswood, C. J., Kibuka, M., & Thornton, J. G. (2013). Position in the second stage of labour for women with epidural anaesthesia. *Cochrane Database of Systematic Reviews*(1), CD008070. <https://doi.org/10.1002/14651858.CD008070.pub2>
- Lawrence, A., Lewis, L., Hofmeyr, G. J., & Styles, C. (2013). Maternal positions and mobility during first stage labour. *Cochrane Database of Systematic Reviews*(8), CD003934. <https://doi.org/10.1002/14651858.CD003934.pub3>
- Mathew, A., Nayak, S., & Vandana, K. (2012). A comparative study on effect of ambulation and birthing ball on maternal and newborn outcome among primigravida mothers in selected hospitals in Mangalore. *Journal of Health and Allied Sciences NU*, 2(02), 2-5. <https://doi.org/10.1055/s-0040-1703561>
- Ministry of Health of Saudi Arabia. (2019). *East Jeddah Hospital statistics*. <https://www.moh.gov.sa/en/Ministry/Statistics/Indicator/Pages/Indicator-1440.aspx>
- Modoor, S., Fouly, H., & Rawas, H. (2021). The effect of warm compresses on perineal tear and pain intensity during the second stage of labor: A randomized controlled trial. *Belitung Nursing Journal*, 7(3), 210-218. <https://doi.org/10.33546/bnj.1452>
- Mselle, L. T., & Eustace, L. (2020). Why do women assume a supine position when giving birth? The perceptions and experiences of postnatal mothers and nurse-midwives in Tanzania. *BMC Pregnancy and Childbirth*, 20(1), 1-10. <https://doi.org/10.1186/s12884-020-2726-4>
- Oats, J. J. N., & Abraham, S. (2017). *Llewellyn-Jones fundamentals of obstetrics and gynaecology E-book* (10th ed.). Elsevier Health Sciences.
- Polit, D. F., & Beck, C. T. (2018). *Essentials of nursing research: Appraising evidence for nursing practice* (9th ed.). Philadelphia:: Wolters Kluwer. <https://books.google.co.th/books?id=Z6DwjwEACA-AJ>
- Priddis, H., Dahlen, H., & Schmied, V. (2012). What are the facilitators, inhibitors, and implications of birth positioning? A review of the literature. *Women and Birth*, 25(3), 100-106. <https://doi.org/10.1016/j.wombi.2011.05.001>
- Ricci, S. S. (2012). *Essentials of maternity, newborn, & women's health nursing* (3rd ed.). Philadelphia: Lippincott Williams & Wilkins.
- Shenbagavalli, S., & Menaka, K. (2016). Effectiveness of selected nursing intervention on maternal and new born outcome among primigravida mothers at Selected Hospitals, Tamilnadu. *Asian Journal of Nursing Education and Research*, 6(3), 407-410. <http://dx.doi.org/10.5958/2349-2996.2016.00072.0>
- Walker, K. F., Kibuka, M., Thornton, J. G., & Jones, N. W. (2018). Maternal position in the second stage of labour for women with epidural anaesthesia. *Cochrane Database of Systematic Reviews*(11), CD008070. <https://doi.org/10.1002/14651858.CD008070.pub4>
- Watson, H. L., & Cooke, A. (2018). What influences women's movement and the use of different positions during labour and birth: A systematic review protocol. *Systematic Reviews*, 7(1), 1-6. <https://doi.org/10.1186/s13643-018-0857-8>
- White, L., Duncan, G., & Baumle, W. (2011). *Foundation of maternal & pediatric nursing* (3rd ed.). Boston: Delmar Cengage Learning.
- Wong-Baker FACES Foundation. (2019). *Wong-Baker FACES® Pain Rating Scale*. <http://www.WongBakerFACES.org>
- World Health Organization. (2018). *WHO recommendations intrapartum care for a positive childbirth experience*. Geneva: World Health Organization.
- Zaky, N. H. (2016). Effect of pelvic rocking exercise using sitting position on birth ball during the first stage of labor on its progress. *IOSR Journal of Nursing*, 5(4), 19-27.

Cite this article as: Al Aryani, Z., Orabi, A., & Fouly, H. (2022). Examining the impact of upright and recumbent positions on labor outcomes in Saudi Arabia: A quasi-experiment. *Belitung Nursing Journal*, 8(4), 316-324. <https://doi.org/10.33546/bnj.2114>